

# Factors Related to Advanced Stage of Cancer Presentation in Botswana

abstract

**Purpose** Botswana, a country with a high prevalence of HIV, has an increasing incidence of cancer-related mortality in the post-antiretroviral therapy era. Despite universal access to free health care, the majority of Botswana patients with cancer present at advanced stages. This study was designed to explore the factors related to advanced-stage cancer presentation in Botswana.

**Methods** Patients attending an oncology clinic between December 2015 and January 2017 at Princess Marina Hospital in Gaborone, Botswana, completed a questionnaire on sociodemographic and clinical factors as well as cancer-related fears, attitudes, beliefs, and stigma. Odds ratios (ORs) were calculated to identify factors significantly associated with advanced stage (stage III and IV) at diagnosis.

**Results** Of 214 patients, 18.7% were men and 81.3% were women. The median age at diagnosis was 46 years, with 71.9% of patients older than 40 years. The most commonly represented cancers included cervical (42.3%), breast (16%), and head and neck (15.5%). Cancer stages represented in the study group included 8.4% at stage I, 19.2% at stage II, 24.1% at stage III, 11.9% at stage IV, and 36.4% at an unknown stage. Patients who presented at advanced stages were significantly more likely to not be afraid of having cancer (OR, 3.48;  $P < .05$ ), believe that their family would not care for them if they needed treatment (OR, 6.35;  $P = .05$ ), and believe that they could not afford to develop cancer (OR, 2.73;  $P < .05$ ). The perception that symptoms were less serious was also significantly related to advanced stage ( $P < .05$ ). Patients with non-female-specific cancers were more likely to present in advanced stages (OR, 5.67;  $P < .05$ ).

**Conclusion** Future cancer mortality reduction efforts should emphasize cancer symptom awareness and early detection through routine cancer screening, as well as increasing the acceptability of care-seeking, especially among male patients.

© 2018 by American Society of Clinical Oncology Licensed under the Creative Commons Attribution 4.0 License

Chidinma Anakwenze

Rohini Bhatia

William Rate

Lame Bakwenabatsile

Kebatsabile Ngoni

Sarah Rayne

Preet Dhillon

Mohan Narasimhamurthy

Ari Ho-Foster

Doreen Ramogola-Masire

Surbhi Grover

Author affiliations and support information (if applicable) appear at the end of this article.

**Corresponding author:**

Surbhi Grover, MD, MPH, Department of Radiation Oncology, University of Pennsylvania, 3400 Civic Center Blvd, Philadelphia, PA 19104; e-mail: [surbhi.grover@uphs.upenn.edu](mailto:surbhi.grover@uphs.upenn.edu).

**INTRODUCTION**

As low- and middle-income countries experience population growth and reduced mortality from communicable diseases, their cancer burdens increase. By 2030, cancer rates will nearly double in some low- and middle-income countries where screening programs are scarce, health systems are poorly equipped, and awareness is limited.<sup>1</sup> Because of these limitations, patients often present with advanced-stage malignancies, which leads to greater rates of cancer-related deaths.<sup>2</sup>

Botswana, a middle-income country in southern Africa with an HIV prevalence of 21.9% among adults 15 to 49 years of age, has an increasing incidence of cancer-related mortality in the post-antiretroviral therapy era.<sup>3</sup> With a population of approximately 2 million people, Botswana has 1,600 new patients with cancer per year.<sup>1</sup> The country's name means land of the

Tswana, referring to the dominant ethnic group in Botswana. However, the term Batswana is used generally as a demonym for all citizens of Botswana.<sup>4</sup>

Cancer screening is not common in the public sector of Botswana. Prior studies have reported low rates of mammography screening, likely because mammography is not readily available and thus is not part of routine screening.<sup>5</sup> Prostate cancer screening is also not routine.<sup>6</sup> Similarly, colon cancer screening is not commonly performed, partly because the incidence of colon cancer in Botswana is low compared with higher-income countries.<sup>1,7</sup> However, efforts to bolster cervical cancer screening have been made; for example, See and Treat, a program involving visual inspection after acetic acid application to the cervix, was implemented for HIV-infected women.<sup>8,9</sup> Additional efforts to curb the incidence of cervical cancer have also included

a government-funded comprehensive human papillomavirus vaccination plan.<sup>10</sup> Despite universal access to government-funded health care, the majority of Botswana patients with cancer, half of whom are infected with HIV, present at advanced stages.<sup>11</sup> Although antiretroviral therapy coverage has reached 83% in Botswana, and median CD4 counts in previously published literature from the antiretroviral therapy era demonstrate a well-managed HIV population, HIV-infected Botswana individuals remain three to five times more likely to develop cancer than age-matched HIV-negative controls.<sup>3,8,12,13</sup> Surprisingly, even HIV-infected patients with cancer with regular longitudinal contact with the health care system do not have faster linkages into cancer care.<sup>11</sup> Median time from cancer symptom onset to treatment initiation in Botswana was reported to be 13 months, compared with 3 months in more developed settings.<sup>11,14,15</sup>

Prior studies suggest that delays in oncologic treatment may be related to distance from the hospital, health insurance status, quality of health care systems, use of traditional healers, financial opportunity costs, limited cancer awareness, and cancer stigma or fear.<sup>16-18</sup> A prospective study conducted in Botswana suggested that compared with patients with early-stage disease, patients with rapidly progressing symptoms and advanced disease entered into specialized oncology treatment earlier after initial symptom recognition.<sup>11</sup> Perhaps additional improvements can be made through earlier symptom recognition at the individual or clinic level. However, additional efforts are needed to understand factors associated with advanced-stage presentation.

Given the existing literature, we suspect that the reason for advanced presentation is multifactorial. Therefore, we sought to describe sociodemographic and clinical factors, as well as the knowledge, attitudes, and beliefs associated with advanced stage at diagnosis in Botswana. We believe that understanding factors associated with advanced-stage presentation is crucial to facilitating earlier cancer detection and intervention, thus reducing cancer-related mortality in Botswana.

## METHODS

### Procedure

This was a cross-sectional study conducted from December 2015 to January 2017 at the Princess Marina Hospital in Gaborone, Botswana. This hospital provides oncology care for the majority of patients in southern Botswana. The study population consisted of a convenience sample of 214 newly diagnosed patients who were at least 18 years old and presented to Princess Marina Hospital for initial cancer treatment with a pathologically confirmed diagnosis of cancer.

Patients were approached by a member of the research team and asked whether they would like to participate in a study assessing delays in cancer care. The questionnaire was administered in Setswana and English, and research assistants administered the survey to patients who were illiterate in the study language. The study protocol was approved by the Institutional Review Board at the University of Pennsylvania and the Health Research Development Committee at the Botswana Ministry of Health. Written informed consent was obtained from each participant before completion of the questionnaire.

### Study Measures

The first section of the questionnaire consisted of sociodemographic questions, including age, sex, relationship status, literacy level, and educational attainment. We also assessed the presence of comorbidities (diabetes, HIV, and tuberculosis), distance to the hospital, symptom severity, and cancer site (Table 1). Additional questions assessed place of residence, languages spoken, employment, economic status, ability to take time off from work, assets (home, land, or livestock ownership), family size, methods of transportation, travel time to the hospital, and religious background (Table 2).

The final section was adapted from a prior study among patients with breast cancer in South Africa. It assessed cancer-related fears, attitudes, beliefs, and stigma using a four-point summative scale ranging from strongly agree to strongly disagree (Table 3).<sup>19</sup> For the analysis, patients who agreed or strongly agreed were categorized as agree and patients who disagreed or strongly disagreed were categorized as disagree. The questionnaire was piloted with Botswana

**Table 1.** Patient Characteristics by Stage

Characteristic	Early Stage (I and II) No. (%)	Late Stage (III and IV) No. (%)	Stage Unknown No. (%)	Response Rate %	Adjusted Probability (P)
Age (years)				100	.6025
20-39	18 (8.9)	19 (9.4)	20 (9.9)		
40-59	28 (13.8)	43 (21.2)	39 (19.2)		
≥ 60	10 (4.9)	11 (5.4)	15 (7.4)		
Sex				100	.0471
Male	2 (0.9)	8 (3.7)	31 (14.5)		
Female	54 (25.2)	65 (30.3)	50 (23.4)		
Relationship status				98.13	.1360
Single	41 (19.5)	40 (19.0)	51 (24.3)		
Married/in a serious relationship	9 (4.3)	24 (11.4)	22 (10.5)		
Living with a partner	3 (1.4)	4 (1.9)	1 (0.5)		
Divorced/separated/widowed	3 (1.4)	6 (2.9)	6 (2.9)		
Literate in own language				99.53	.5909
Yes	47 (22.1)	61 (28.6)	64 (30.0)		
No	9 (4.2)	15 (7.0)	17 (8.0)		
Literate in English				98.60	.7785
Yes	24 (11.4)	34 (16.1)	40 (19.0)		
No	32 (15.2)	41 (19.4)	40 (19.0)		
Education level				81.78	.6193
No formal education	6 (3.4)	11 (6.3)	11 (6.3)		
Primary school only	20 (11.4)	21 (12.0)	18 (10.3)		
Secondary school	19 (10.9)	29 (16.6)	22 (12.6)		
Tertiary/postbachelor's degree	4 (2.3)	6 (3.4)	6 (3.4)		
Other	0 (0.0)	2 (1.1)	0 (0.0)		
Diabetic				99.53	.2859
Yes	2 (0.9)	6 (2.8)	2 (0.9)		
No	55 (25.8)	69 (32.4)	79 (37.1)		
HIV status				98.60	.4094
Positive	38 (18.0)	45 (21.3)	45 (21.3)		
Negative	18 (8.5)	29 (13.7)	36 (17.1)		
Hypertensive				99.53	.7204
Yes	12 (5.6)	18 (8.5)	17 (8.0)		
No	45 (21.1)	58 (27.2)	63 (29.6)		
History of tuberculosis				98.60	.5422
Yes	3 (1.4)	6 (2.8)	15 (7.1)		
No	53 (25.1)	68 (32.2)	66 (31.3)		
Distance to hospital (km)				96.26	.0733
5-50	15 (7.3)	29 (14.1)	23 (11.2)		
51-200	20 (9.7)	19 (9.2)	22 (10.7)		
201-400	12 (5.8)	7 (3.4)	15 (7.3)		
≥ 400	9 (4.4)	19 (9.2)	15 (7.3)		
Unable to locate village	0 (0.0)	0 (0.0)	1 (0.5)		

(Continued on following page)

**Table 1.** Patient Characteristics by Stage (Continued)

Characteristic	Early Stage (I and II) No. (%)	Late Stage (III and IV) No. (%)	Stage Unknown No. (%)	Response Rate %	Adjusted Probability (P)
Severity of symptoms				97.66	.0428
Not serious	10 (4.8)	17 (8.1)	22 (10.5)		
A little serious	14 (6.7)	14 (6.7)	11 (5.3)		
Moderately serious	3 (1.4)	11 (5.3)	9 (4.3)		
Serious	10 (4.8)	3 (1.4)	8 (3.8)		
Very serious	19 (9.1)	29 (13.9)	29 (13.9)		
Cancer diagnosis site				99.53	.0523
Cervical	38 (17.8)	38 (17.8)	14 (6.6)		
Breast	10 (4.7)	19 (8.9)	5 (2.3)		
Head and neck	0 (0.0)	10 (4.7)	23 (10.8)		
Vulvar	6 (2.8)	5 (2.3)	4 (1.9)		
Kaposi sarcoma	0 (0.0)	1 (0.5)	13 (6.1)		
Endometrial	1 (0.5)	1 (0.5)	5 (2.3)		
Penile	0 (0.0)	2 (0.9)	4 (1.9)		
Anal	1 (0.5)	0 (0.0)	4 (1.9)		
Esophageal	0 (0.0)	0 (0.0)	5 (2.3)		
Lymphoma	0 (0.0)	0 (0.0)	3 (1.4)		
Prostate	1 (0.5)	0 (0.0)	0 (0.0)		

women to further refine it and adapt it to Botswana culture.

Cancer diagnosis was made using pathologic confirmation. Cancers were staged according to the TNM staging system of the American Joint Commission on Cancer, seventh edition.<sup>20</sup> Staging evaluation varied by cancer site but often included medical chart review, physical examination, and imaging with chest x-ray and ultrasound. Cancers were categorized as early (stage I and II) or advanced (stage III and IV). However, some patients were characterized as being in the unknown stage if they were not staged before treatment initiation or if imaging modalities were not functional at the time of diagnosis.

### Statistical Analyses

Responses to the questionnaire were collected electronically using REDCap (Research Electronic Data Capture) tools hosted at the University of Pennsylvania.<sup>21</sup> All statistical analysis was completed using commercially available analytic software (STATA, version 15.0; STATA, College Station, TX).

Nonparametric post hoc one-way analysis of variance between all measured and calculated variables for patients with early-stage disease

(stage I and II) and advanced-stage disease (stage III and IV) were analyzed using the Kruskal-Wallis H test, which adjusts for ties between ordinal responses to a question. Unadjusted and adjusted probabilities were reported to indicate the significance of the difference between recorded categorical patient responses in the early- and advanced-stage groups (Table 3). Univariable logistic regression was used to determine directionality of the difference between early- and advanced-stage group responses to the fears, attitudes, beliefs, and stigma portion of the survey, as well as unadjusted odds ratios (ORs) describing the impact of our population's agreement or disagreement with survey questions on their relative probability of being in the advanced-stage group (Tables 3 and 4).

A multivariable logistic regression analysis was performed for each variable contained in the surveys to determine the significance of any single item's ability to predict advanced stage at diagnosis in a post hoc fashion. The significance of both the model and the categorical factors is listed for all items producing a significant predictive model in Table 4, with associated ORs. Any categorical factors with an SE of greater than 1,000 were excluded from the logistic regression performed on a variable.

**Table 2.** Sociodemographic and Economic Factors by Stage

Factor	All Stages No. (%)	Early Stage (I and II) No. (%)	Late Stage (III and IV) No. (%)	Response Rate %	Adjusted Probability <i>P</i>
Place of residence				98.50	.3946
Village	83 (63.4)	38 (29.0)	45 (34.4)		
Town	36 (27.5)	15 (11.5)	21 (16.0)		
City	12 (9.2)	3 (2.3)	9 (6.9)		
Total	131	56 (42.7)	75 (57.3)		
Languages spoken by patients				100	
Setswana	129 (97.0)	57 (42.9)	72 (54.1)		.0798
English	60 (45.1)	24 (18.0)	36 (27.1)		.5476
Sekalaka	9 (6.8)	5 (3.8)	4 (3.0)		.4270
Sekgalagadi	4 (3.0)	3 (2.3)	1 (0.8)		.1888
Afrikaans/Sesubiya/ Sesarwa	4 (3.0)	1 (0.8)	3 (2.3)		.2189/.2482/.3865
Employment status at diagnosis				100	.7542
Employed	47 (35.3)	21 (15.8)	26 (19.5)		
Unemployed	86 (64.7)	36 (27.1)	50 (37.6)		
Total	133				
Monthly salaries of those employed at diagnosis				87.23	.1582
< P2,000	4 (8.5)	0 (0.0)	4 (8.5)		
P2,001-5,000	34 (72.3)	17 (36.2)	17 (36.2)		
> P5,001	3 (6.4)	1 (2.1)	2 (4.3)		
Not reported	6 (12.8)	3 (6.4)	3 (6.4)		
Difficulty taking time off from work				97.87	.3787
Always	5 (10.6)	2 (4.3)	3 (6.4)		
Mostly	2 (4.3)	1 (2.1)	1 (2.1)		
Sometimes	5 (10.6)	4 (8.5)	1 (2.1)		
Not really	34 (72.3)	13 (27.7)	21 (44.7)		
Not reported	1 (2.1)	1 (2.1)	0 (0.0)		
Monthly family income					.5729
< P3,000	17 (7.9)	9 (6.8)	8 (6.0)		
P3,001-6,000	65 (30.4)	26 (19.5)	39 (29.3)		
> P6,001	6 (2.8)	2 (1.5)	4 (3.0)		
Not reported	45 (21.0)	20 (15.0)	25 (18.8)		
Home ownership				97.74	1.0000
Owns a home	72 (54.1)	31 (23.3)	41 (30.8)		
Does not own a home	58 (43.6)	25 (18.8)	33 (24.8)		
Not reported	3 (2.3)	1 (0.8)	2 (1.5)		
Land ownership				96.24	.4315
Owns land	52 (39.1)	19 (14.3)	33 (24.8)		
Does not own land	76 (57.1)	35 (26.3)	41 (30.8)		
Not reported	5 (3.8)	3 (2.3)	2 (1.5)		

(Continued on following page)

**Table 2.** Sociodemographic and Economic Factors by Stage (Continued)

Factor	All Stages No. (%)	Early Stage (I and II) No. (%)	Late Stage (III and IV) No. (%)	Response Rate %	Adjusted Probability <i>P</i>
Livestock ownership				94.74	.4344
Owns livestock	29 (21.8)	9 (6.8)	20 (15.0)		
Does not own livestock	97 (72.9)	44 (33.1)	53 (39.8)		
Not reported	7 (5.3)	4 (3.0)	3 (2.3)		
No. of family members				96.24	.9310
0-1	9 (6.8)	4 (3.0)	5 (3.8)		
2-3	40 (30.1)	16 (12.0)	24 (18.0)		
4-5	34 (25.6)	14 (10.5)	20 (15.0)		
> 5	45 (33.8)	21 (15.8)	24 (18.0)		
Not reported	5 (3.8)	2 (1.5)	3 (2.3)		
No. of dependents				93.23	.2474
0-1	45 (33.8)	24 (18.0)	21 (15.8)		
2-3	51 (38.3)	21 (15.8)	30 (22.6)		
4-5	19 (14.3)	5 (3.8)	14 (10.5)		
> 5	9 (6.8)	4 (3.0)	5 (3.8)		
Not reported	9 (6.8)	3 (2.3)	6 (4.5)		
Most common method of transportation				95.49	.6428
Taxi	32 (24.1)	11 (8.3)	21 (15.8)		
Bus	75 (56.4)	34 (25.6)	41 (30.8)		
Personal or family car	19 (14.3)	9 (6.8)	10 (7.5)		
Borrow a car or get a ride	1 (0.8)	0 (0.0)	1 (0.8)		
Not reported/other	6 (4.5)	3 (2.3)	3 (2.3)		
Ease of transportation to the hospital				96.99	.9937
I am independent	88 (66.2)	38 (28.6)	50 (37.6)		
I am dependent on others	34 (25.6)	14 (10.5)	20 (15.0)		
Getting to the hospital can be burdensome	7 (5.3)	3 (2.3)	4 (3.0)		
Not reported/other	4 (3.0)	2 (1.5)	2 (1.5)		
Travel time to the hospital				99.25	.5916
< 30 minutes	5 (3.8)	1 (0.8)	4 (3.0)		
30 minutes to 1 hour	51 (38.3)	23 (17.3)	28 (21.1)		
1 hour to 4 hours	48 (36.1)	22 (16.5)	26 (19.5)		
> 4 hours	28 (21.1)	10 (7.5)	18 (13.5)		
Not reported	1 (0.8)	1 (0.8)	0 (0.0)		
Religious beliefs				96.24	.0569
Christian	111 (51.9)	53 (39.8)	58 (43.6)		
Muslim	1 (0.5)	0 (0.0)	1 (0.8)		

(Continued on following page)

**Table 2.** Sociodemographic and Economic Factors by Stage (Continued)

Factor	All Stages No. (%)	Early Stage (I and II) No. (%)	Late Stage (III and IV) No. (%)	Response Rate %	Adjusted Probability <i>P</i>
Traditional African	2 (0.9)	0 (0.0)	2 (1.5)		
Believe in a god or higher power but not religious	10 (4.7)	1 (0.8)	9 (6.8)		
Do not believe in a god or higher power	4 (1.9)	1 (0.8)	3 (2.3)		
Not reported/other	5 (2.3)	2 (1.5)	3 (2.3)		

Abbreviation: P, Botswana pula.

## RESULTS

### Demographics

A total of 214 patients presenting for specialized cancer care were included in the analysis. Of the 214 patients, 57 (28.2%) were younger than 40 years of age, 40 (18.7%) were male, and 174 (81.3%) were female. Half of the patients (50.3%) reported at least a secondary school level of education, and 55 (26.2%) reported being married or living with a partner. The most commonly represented cancers were cervical ( $n = 90$ ; 42.3%), breast ( $n = 34$ ; 16%), head and neck ( $n = 33$ ; 15.5%), vulvar ( $n = 15$ ; 7%), and Kaposi sarcoma ( $n = 14$ ; 6.6%). The remaining 27 malignancies (12.7%) included endometrial, penile, anal, esophageal, lymphoma, and prostate. At least 49% of the patients with cervical cancer, 11% of the patients with breast cancer, 11% of the patients with head and neck cancer, and 9.5% of the patients with vulvar cancer and Kaposi sarcoma were HIV positive. All other cancers had less than 5% of patients who were HIV positive. Cancer stages represented in the study group included 17 (8.4%) at stage I, 39 (19.2%) at stage II, 49 (24.1%) at stage III, 24 (11.9%) at stage IV, and 74 (36.4%) with unknown stage. Other key patient characteristics are listed by stage in [Table 1](#).

### Socioeconomic Factors

Multiple languages were represented in the patient sample, with 129 patients (97%) speaking Setswana and 60 (45.1%) speaking English, normally as a second language. Most patients reported no transportation problems, with only 17 patients (12.8%) with early-stage cancer and 24 patients (18%) with advanced-stage cancer reporting that getting to the hospital was

burdensome or that they were dependent on others to get to the hospital. Most of the patients ( $n = 86$ ; 64.7%) who responded to the employment question were unemployed at the time of diagnosis. Of those who were employed, 34 (72.3%) earned in Botswana pula (P) between P2,001 and P5,000 per month (\$209 to \$522 in US dollars in June 2016). Other socioeconomic factors are listed by stage in [Table 2](#). No significant relationships were noted between any socioeconomic variable and advanced-stage disease.

### Fears, Attitudes, Beliefs, and Stigma

Patients who were not afraid of having cancer were more likely to present with advanced-stage cancer at diagnosis (OR, 3.48;  $P < .05$ ). Patients who agreed that their family would not care for them if they needed treatment were more likely to have advanced-stage disease (OR, 6.35;  $P = .05$ ). Patients who agreed that they could not afford to develop cancer were more likely to present with advanced disease (OR, 2.73;  $P < .05$ ). Other fears, attitudes, beliefs, and stigmas are listed by stage in [Table 3](#).

### Factors Associated With Advanced Stage in Multivariable Logistic Analysis

Patients with non–female-specific cancers were more likely to present with advanced-stage disease (OR, 5.67;  $P < .05$ ). Female-specific cancers were defined as cancers of the female reproductive organs (cervical, ovarian, uterine, vaginal, vulvar) and breast. There was also a nonsignificant trend toward male patients presenting with advanced-stage cancer (OR, 6.59;  $P = .08$ ; [Table 4](#)). Patients with cervical and vulvar cancer were less likely to present with advanced

**Table 3.** Fears, Attitudes, Beliefs, and Stigma by Cancer Stage

Fears, Attitudes, Beliefs, and Stigma	Unadjusted Probability (P)	Adjusted Probability (P)	Responses	Stage I/II Agree Responses	Stage III/IV Agree Responses (OR)
<b>Fears</b>					
Scared of having cancer	<b>.0368</b>	<b>.0148</b>	<b>131</b>	<b>13</b>	<b>6 (0.287)</b>
Scared of job loss	.1927	.1276	107	7	4 (0.374)
Scared of surgery	.6518	.5977	126	5	5 (0.706)
Scared of missing appointments or treatment due to work	.6853	.6367	116	4	7 (1.36)
Scared of missing appointments or treatment due to transport problems	.6500	.5961	132	7	12 (1.31)
Scared of radiation therapy	.8069	.7753	132	6	7 (0.845)
Scared of chemotherapy side effects	.4228	.3491	132	9	8 (0.614)
Scared of dying from cancer	.3906	.3159	132	11	10 (0.619)
<b>Attitudes</b>					
My family will not care for me if I need treatment	.0977	.0527	131	1	8 (6.35)
I have no family history of cancer, so I should not have gotten cancer	.4446	.3715	128	17	18 (1.410)
I can beat cancer	.7937	.7597	131	54	74 (0.632)
I can't afford to get cancer	<b>.0683</b>	<b>.0333</b>	<b>129</b>	<b>32</b>	<b>29 (2.73)</b>
People with cancer can survive	.9795	.9760	130	52	71 (0.687)
<b>Beliefs</b>					
Old people get cancer	.6926	.6434	127	28	36 (0.386)
Cancer is a curse and the result of past sins or bewitchment	.6148	.5574	130	5	9 (0.340)
Using contraceptive pills or injections can cause cancer	<b>.0487</b>	<b>.0211</b>	<b>129</b>	<b>19</b>	<b>13 (0.723)</b>
Using hormone replacement pills after menopause can cause cancer	<b>.0360</b>	<b>.0142</b>	<b>128</b>	<b>17</b>	<b>10 (0.955)</b>
My family will support me	.5291	.4616	131	53	71 (1.39)
My faith in God will heal me	.5753	.5120	129	43	56 (2.23)
Getting cancer is part of God's plan	.4496	.3755	128	32	51 (1.55)
If someone gets cancer, it doesn't matter when they find out, they will still die	.2563	.1851	129	4	11 (0.798)
Cancer kills most people who get it	.6876	.6391	128	10	11 (0.464)
People have been cured using traditional nonmedical healing	.8261	.7971	129	3	5 (1.05)
People have been cured through prayer and faith alone	.1038	.0566	128	41	46 (0.704)
Cancer can be treated in many ways, not just through medicine and surgery	.9170	.9026	127	23	34 (0.560)
People with cancer will die when they go to the hospital	.6817	.6313	129	4	4 (1.02)
There are treatments for cancer	.4846	.4141	130	52	68 (0.677)
People have been cured of cancer using alternative therapies	.3637	.2883	128	24	26 (0.322)
Early detection and timely treatment will result in a longer life	.2143	.1461	129	52	67 (1.45)
Some cancer treatments could kill me	.5577	.4941	128	6	11 (1.56)
If someone gets cancer, a lot of different treatments won't make any difference	.4932	.4242	128	5	10 (0.769)

(Continued on following page)



**Table 3.** Fears, Attitudes, Beliefs, and Stigma by Cancer Stage (Continued)

Fears, Attitudes, Beliefs, and Stigma	Unadjusted Probability ( <i>P</i> )	Adjusted Probability ( <i>P</i> )	Responses	Stage I/II Agree Responses	Stage III/IV Agree Responses (OR)
Stigma					
It is my fault that I have a health condition	.2867	.2128	126	4	2 (0.345)
Family members or neighbors visit less or stopped visiting due to stigma	.7965	.7628	129	1	2 (1.45)
My relationship with my spouse has changed or worsened due to stigma	.4518	.3785	110	2	1 (0.350)
I have been kept isolated in my home due to stigma	.5441	.4784	130	1	3 (2.25)
People have prevented me from sharing food due to stigma	.5517	.4862	131	1	3 (2.22)
People have prevented me from playing with their children due to stigma	.3150	.2398	131	1	0 < (0.001)
Family members refused to sleep in the same room as me due to stigma	.5621	.4970	130	1	3 (2.18)
Family members or neighbors have spoken badly of me due to stigma	.6977	.6499	130	2	4 (1.49)
People have blamed me for my health problems	.9377	.9271	131	2	3 (1.09)

NOTE. Bold type indicates significant adjusted *P* values.

Abbreviation: OR, odds ratio.

stages (OR, 0.07 and OR, 0.06, respectively;  $P < .05$ ). Patients who reported serious symptom severity at presentation were less likely to have advanced-stage disease (OR, 0.176;  $P < .05$ ).

## DISCUSSION

Half of the study participants at the Princess Marina Hospital presented with advanced-stage cancers. Understanding factors associated with advanced-stage presentation is crucial to facilitating earlier cancer detection and intervention, thus reducing cancer-related mortality. This is particularly important in cancers that are preventable through screening and vaccination, such as cervical cancer, human papillomavirus-related cancers, and breast cancer, which typically present at an advanced stage in Botswana and comprise more than half of the study population when combined.

Our study population is representative of the general population when considering age, sex, socioeconomic status, and cancer type. For example, the median age at diagnosis was 46 years, with 71.9% of patients older than 40 years. This is consistent with Botswana National Cancer Registry data reporting that the median age at cancer diagnosis was 47 years for women and 50 years for men.<sup>22</sup> Our study also showed

that 72.3% of the participants earned between P2,001 and P5,000 per month, which is within the limits of the average monthly salary of Botswana (P4,801) during the study period.<sup>23</sup> The most commonly represented cancers included cervical (42.3%) and breast (16%). These data are consistent with existing cancer incidence data, which show that cervical and breast cancer are the most commonly diagnosed cancers, as well as the most common causes of cancer death, representing 15.3% and 9.5%, respectively, of all newly diagnosed cancers. However, given the large number of female participants in our study, cervical and breast cancers had higher representation. Patients with Kaposi sarcoma comprised 6.6% of our sample, which is consistent with existing data that show Kaposi sarcoma represents 7.6% of all newly diagnosed cancers.<sup>1</sup>

There is relatively good access to specialized cancer care in Botswana, with government-funded care being accessible to 90% of the population.<sup>8</sup> Although patients with HIV might have closer linkages to the health care system through Botswana's robust antiretroviral therapy program, our study did not show an association between HIV status and earlier stage at diagnosis. This is consistent with prior data that suggest that even HIV-infected patients with cancer

**Table 4.** Logistic Regression Analysis for Variables Predicting Advanced Stage of Presentation

Variable	Model Significance (P)	Factor Significance (P)	Odds Ratio
Gender	.0313		
Male		.08	6.59
Cancer diagnosis site <sup>a</sup>	.0057		
Cervical		.013	0.0714
Vulvar		.019	0.0595
Breast		.071	0.136
Anal		.990	< 0.0001
Prostate		.990	< 0.0001
Symptom severity	.0361		
Not serious		.828	1.11
A little serious		.378	0.588
Moderately serious		.220	2.16
Serious		.024	0.176
Very serious		.828	0.898
Predominantly female cancer	.0089		
Female cancer		.026	0.176
Nonfemale cancer		.026	5.67

NOTE. Model significance *P* values reflect  $\chi^2$  differences between early-stage and advanced-stage groups for each.

<sup>a</sup>Excluded head and neck plus penile because of SE > 1,000; endometrial used as base.

with regular contact with the health care system do not have faster linkages into cancer care.<sup>11</sup> This also suggests that there is a need to indiscriminately bolster existing methods for early diagnosis, provision of quality care, and efficient management of limited resources in patients with HIV-positive and HIV-negative disease. It is important to note that as antiretroviral therapy coverage has increased from 7.3% to 82.3% between 2003 and 2008, age-adjusted cancer incidence has decreased in patients with HIV by 8.3% per year. However, with a progressively larger and older HIV population, there is still a high number of incident cancers in the HIV population.<sup>22</sup>

Prior studies have suggested that advanced-stage presentation may be related to difficulty getting to the hospital; however, our study found that difficulty getting to the hospital was not a cause of advanced-stage presentation.<sup>16,18</sup> This suggests that there may be other prevailing causes in Botswana. For example, gender-related factors predicted increased stage at presentation. A qualitative meta-analysis reported that men often view help-seeking as unmasculine, and women

find help-seeking easier, given their greater contacts with health services for themselves and their families.<sup>18</sup> One study reported that of 5,000 monthly patients attending a clinic in Gaborone, Botswana, 60% were female. Women were seen more frequently than men for reasons such as sexual and reproductive health, pre-employment medical examination, and consultation for various new symptoms.<sup>5</sup> The discrepancies between male and female stage of presentation may also be related to differences in symptoms between gender-specific cancers (ie, vaginal bleeding may be more disturbing than urinary symptoms in prostate cancer).

In this study, patients who perceived their symptoms to be serious were more likely to present at early stages. This is consistent with prior studies that suggest that severe symptoms, symptoms that interfere with everyday life, or well-recognized symptoms (ie, lump) are potent triggers of early illness recognition and help-seeking.<sup>18,24</sup> In contrast, patients with vague or nonspecific initial symptoms are known to have delayed illness recognition.

We also found that certain fears, attitudes, and beliefs were predictive of advanced stage at presentation. Patients who were not afraid of having cancer were more likely to present with advanced-stage disease. This suggests that those who present with advanced-stage cancer may not understand the mortality risks associated with a cancer diagnosis or may not be concerned about this within their belief structures. This is a direct contradiction of the other highly prevalent belief that early detection and early treatment results in a longer life. These contradicting beliefs represent a cognitive dissonance that aids in the avoidance of their own mortality, particularly in advanced-stage cancer. Most of the participants also felt that their faith would cure their disease. This belief in faith-based cure has been previously reported in other African countries.<sup>19</sup> Patients who believed that their family would not care for them or that they could not afford to develop cancer were more likely to present with advanced-stage disease. This is consistent with prior studies that suggest responsibility for the needs of other family members often prevents patients from prioritizing their own health needs.<sup>18</sup> Often, patients do not seek health care until their symptoms start to affect their ability to work.<sup>25,26</sup> These findings suggest

that future interventions should target increasing cancer symptom awareness and promotion of early care-seeking among primary wage earners in the family.

Most of the patients with early- and advanced-stage disease held appropriate fears, attitudes, and beliefs regarding cancer and its treatment options. They also indiscriminately reported low levels of cancer stigmatization in their homes and communities, which is consistent with prior African studies that report low levels of cancer stigma.<sup>27-29</sup> This is important to note because stigma has been previously reported as a potential barrier to participation in cancer screening or cancer care-seeking activities.<sup>30</sup> Although interventions targeted toward reducing stigma might improve the psychosocial well-being, interpersonal relationships, and financial opportunities for patients with cancer and can even increase early health care-seeking behaviors, the lack of stigma experienced in this patient population suggests that an emphasis on stigma reduction may not play an important role in the Botswana population of patients with cancer.<sup>31,32</sup>

This study is not without limitations. The sample size was small, which may affect the reliability of the survey's results. Individuals with early-stage cancer may not know that they are sick, and thus may not be well represented in the study sample. Furthermore, we were only able to capture patients who present for care. In addition, self-reported fear, attitude, belief, and stigma data might be biased because of social desirability and recall bias after diagnosis. Issues with survey-item nonresponse may also introduce

bias; however, the majority of survey items had a greater than 90% response rate. We do not have data on patients who refused to participate in the study. Patient refusal could present as a study limitation; however, it is unclear in which direction this might bias the study. Last, the heterogeneity of the cancer sites makes it difficult to form cancer site-specific conclusions that may reduce advanced-stage cancer presentation in the cancer site of interest. Regardless, these findings are important and provide significant insight into causes of advanced-stage cancer presentation in Botswana.

This study examined the sociodemographic and clinical factors, as well as the knowledge, attitudes, and beliefs, associated with delayed stage at diagnosis in Botswana. Patients who presented at advanced stages were more likely to not be afraid of having cancer, believe that their family would not care for them if they needed treatment, and believe that they could not afford to develop cancer. Advanced stage at presentation was found to be associated with non-female-specific cancers and the perception that symptoms were less serious. Future cancer mortality reduction efforts should emphasize cancer symptom awareness and early detection through routine cancer screening, as well as increasing the acceptability of care-seeking through education about cancer outcomes if detected early, especially among male patients.

DOI: <https://doi.org/10.1200/JGO.18.00129>

Published online on [jgo.org](http://jgo.org) on December 11, 2018.

#### AUTHOR CONTRIBUTIONS

**Conception and design:** William Rate, Kebatshabile Ngoni, Mohan Narasimhamurthy

**Administrative support:** Lame Bakwenabatsile

**Provision of study materials or patients:** Lame Bakwenabatsile, Kebatshabile Ngoni

**Collection and assembly of data:** Rohini Bhatia, William Rate, Lame Bakwenabatsile

**Data analysis and interpretation:** Chidinma Anakwenze, Rohini Bhatia, William Rate, Sarah Rayne, Preet Dhillon, Ari Ho-Foster, Doreen Ramogola-Masire, Surbhi Grover

**Manuscript writing:** All authors

**Final approval of manuscript:** All authors

**Accountable for all aspects of the work:** All authors

#### AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to [www.asco.org/rwc](http://www.asco.org/rwc) or [ascopubs.org/jco/site/ifc](http://ascopubs.org/jco/site/ifc).

##### Chidinma Anakwenze

No relationship to disclose

##### Rohini Bhatia

No relationship to disclose

##### William Rate

No relationship to disclose

**Lame Bakwenabatsile**  
No relationship to disclose

**Kebatshabile Ngoni**  
No relationship to disclose

**Sarah Rayne**  
Honoraria: Sysmex

**Preet Dhillon**  
Leadership: Subka Dentist (I)

**Mohan Narasimhamurthy**  
No relationship to disclose

**Ari Ho-Foster**  
No relationship to disclose

**Doreen Ramogola-Masire**  
Travel, Accommodations, Expenses: MSD  
Other Relationship: MSD

**Surbhi Grover**  
No relationship to disclose

#### ACKNOWLEDGMENT

We thank the Department of Radiation Oncology, University of Pennsylvania, for funding; Suneeta Krishnan, Botswana-University of Pennsylvania Partnership; Princess Marina Hospital; and all our patients with cancer.

#### Affiliations

**Chidinma Anakwenze**, MD Anderson Cancer Center, Houston, TX; **Rohini Bhatia**, University of Rochester School of Medicine and Dentistry, Rochester, NY; **William Rate**, Georgetown University School of Medicine, Washington, DC; **Lame Bakwenabatsile**, **Ari Ho-Foster**, and **Surbhi Grover**, Botswana-University of Pennsylvania Partnership, Gaborone, Botswana, and Philadelphia, PA; **Kebatshabile Ngoni**, **Mohan Narasimhamurthy**, and **Doreen Ramogola-Masire**, University of Botswana; **Surbhi Grover**, Princess Marina Hospital, Gaborone, Botswana; **Sarah Rayne**, University of the Witwatersrand, Johannesburg, South Africa; **Preet Dhillon**, Public Health Foundation of India, Gurgaon, Haryana, India; and **Ari Ho-Foster**, **Doreen Ramogola-Masire** and **Surbhi Grover**, University of Pennsylvania, Philadelphia, PA.

#### Support

Funding was provided by the University of Pennsylvania.

#### REFERENCES

1. Ferlay J, Soerjomataram I, Ervik M, et al (eds): GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11. [Internet]. Lyon, France, International Agency for Research on Cancer, 2013
2. Sankaranarayanan R, Swaminathan R, Brenner H, et al: Cancer survival in Africa, Asia, and Central America: A population-based study. *Lancet Oncol* 11:165-173, 2010
3. UNAIDS: Country factsheets, Botswana, <http://www.unaids.org/en/regionscountries/countries/botswana2017>
4. Bolaane M, Mgadla PT: Botswana. New York, NY, Rosen Publishing Group, 1997
5. Tieng'O JG, Pengpid S, Skaal L, et al: Knowledge, attitude and practice of breast cancer examination among women attending a health facility in Gaborone, Botswana. *Gend Behav* 9:3513-3527, 2011
6. Jalloh M, Niang L, Ndoye M, et al: Prostate cancer in Sub Saharan Africa. *J Nephrol Urol Res* 1:15-20, 2013
7. Johnson RH: The cases of cancer seen at a Botswana hospital 1968-1972. *Cent Afr J Med* 21:260-264, 1975
8. Suneja G, Ramogola-Masire D, Medhin HG, et al: Cancer in Botswana: Resources and opportunities. *Lancet Oncol* 14:e290-e291, 2013
9. Ramogola-Masire D, de Klerk R, Monare B, et al: Cervical cancer prevention in HIV-infected women using the "see and treat" approach in Botswana. *J Acquir Immune Defic Syndr* 59:308-313, 2012
10. Raesima MM, Forhan SE, Voetsch AC, et al: Human papillomavirus vaccination coverage among school girls in a demonstration project-Botswana, 2013. *MMWR Morb Mortal Wkly Rep* 64:1147-1149, 2015
11. Brown CA, Suneja G, Tapela N, et al: Predictors of timely access of oncology services and advanced-stage cancer in an HIV-endemic setting. *Oncologist* 21:731-738, 2016

12. Botswana MoH: Botswana National Cancer Registry 2001-2010. Gaborone, Botswana, 2010
13. Grover S, Bvochora-Nsingo M, Yeager A, et al: Impact of human immunodeficiency virus infection on survival and acute toxicities from chemoradiation therapy for cervical cancer patients in a limited-resource setting. *Int J Radiat Oncol Biol Phys* 101:201-210, 2018
14. Hansen RP, Vedsted P, Sokolowski I, et al: Time intervals from first symptom to treatment of cancer: A cohort study of 2,212 newly diagnosed cancer patients. *BMC Health Serv Res* 11:284, 2011
15. Robinson D, Massey T, Davies E, et al: Waiting times for radiotherapy: Variation over time and between cancer networks in southeast England. *Br J Cancer* 92:1201-1208, 2005
16. Jones AP, Haynes R, Sauerzapf V, et al: Travel time to hospital and treatment for breast, colon, rectum, lung, ovary and prostate cancer. *Eur J Cancer* 44:992-999, 2008
17. Langenbach MR, Schmidt J, Neumann J, et al: Delay in treatment of colorectal cancer: Multifactorial problem. *World J Surg* 27:304-308, 2003
18. Smith LK, Pope C, Botha JL: Patients' help-seeking experiences and delay in cancer presentation: A qualitative synthesis. *Lancet* 366:825-831, 2005
19. Rayne S, Schnippel K, Benn C, et al: The effect of access to information on beliefs surrounding breast cancer in South Africa. *J Cancer Educ* 33:806-813, 2018
20. Edge S, Byrd DR, Compton CC, et al (eds): *AJCC Cancer Staging Handbook*. New York, NY, Springer, 2010
21. Harris PA, Taylor R, Thielke R, et al: Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 42:377-381, 2009
22. Dryden-Peterson S, Medhin H, Keabonye-Pusoentsi M, et al: Cancer incidence following expansion of HIV treatment in Botswana. *PLoS One* 10:e0135602, 2015 [Erratum: *PLoS One* 10:e0138742, 2015]
23. Trading Economics ; Botswana Wages: <https://tradingeconomics.com/botswana/wages>
24. Macdonald S, Macleod U, Campbell N, et al: Systematic review of factors influencing patient and practitioner delay in diagnosis of upper gastrointestinal cancer. *Brit J Canc* 94:1272-1280, 2006
25. Burgess C, Hunter MS, Ramirez AJ: A qualitative study of delay among women reporting symptoms of breast cancer. *Br J Gen Pract* 51:967-971, 2001
26. Vaartio H, Kiviniemi K, Suominen T: Men's experiences and their resources from cancer diagnosis to recovery. *Eur J Oncol Nurs* 7:182-190, 2003
27. Ohaeri JU, Campbell OB, Ilesanmi AO, et al: Psychosocial concerns of Nigerian women with breast and cervical cancer. *Psychooncology* 7:494-501, 1998
28. Ohaeri JU, Campbell OB, Ilesanmi AO, et al: The psychosocial burden of caring for some Nigerian women with breast cancer and cervical cancer. *Soc Sci Med* 49:1541-1549, 1999
29. Fort VK, Makin MS, Siegler AJ, et al: Barriers to cervical cancer screening in Mulanje, Malawi: A qualitative study. *Patient Prefer Adherence* 5:125-131, 2011
30. Sankaranarayanan R, Ramadas K, Frie KG, et al: Challenges for Breast and Gynecological Cancer Control by Early Detection in Less-Developed Countries, *Breast and Gynecological Cancers*. New York, NY, Springer, 2013
31. Tod AM, Craven J, Allmark P: Diagnostic delay in lung cancer: A qualitative study. *J Adv Nurs* 61:336-343, 2008
32. Michielutte R, Dignan MB, Sharp PC, et al: Skin cancer prevention and early detection practices in a sample of rural women. *Prev Med* 25:673-683, 1996